

# Health Solutions

The Knowledge for Health

HOME :: June 24, 2010

HowTo

Research

Protocols

eBooks

eLibrary

Videos

Products

Case Studies

Contact

## Favorite Blogs

Nicotine Detox

PEMF-Education

Oxygen Multistep

HowTo Blog

## Sponsors

PEMF Systems

Health Solutions

## Legal Notices

## Bloodroot

### [Printer Friendly PDF](#)

[Sanguinarine is a well known biological agent.](#) It is classified as an escharotic because in high concentration it triggers apoptotic or cell death. Low concentrations selectively disable metabolic compromised cells.

[The video footage](#) suggests that it provides Immunomodulating/directory effects exhibiting elevated activity on pathogenic organism in blood with no apparent damage to white or red blood cells.

[Click Here for current PubMed/NIH research on Sanguinarine](#)

This table suggests a dosing schedule for oral sanguinarine. The schedule is for 5 days on and 2 days off. The 2 days off aid the body to keep up with the detoxification load which often accompanies sanguinarine usage.

1/5 gram is approximately one capsule.

Supplement	Wake	Break	Lunch	Dinner	Bed	Unit	Purpose
<a href="#">Turmeric</a>		1	1	1			Aids absorption and prevents digestive discomfort from Sanguinarine.
<a href="#">Sanguinarine (First 1-3 Months)</a>		2	2	2		Capsule	Empty stomach to aid systemic toxicity by acting outside digestive system.
Acute Care Stage							
<a href="#">Sanguinarine Ongoing (About 4 years)</a>		1	1	1		Capsule	Maintain anti-pathogenic process for a long time until cell lines fully resolved.

Oral dosing uses low systemic doses over time to target systemic pathogens.

Copyright 2005-2009, All Rights Reserved, Mark Squibb